



Target Species

Based on 13,660 complete checklists

289 species observed in North Dakota, US (region/US-ND) in **Jul** that you need for your **World Life List**

| | |
|--------------------------|------------------|
| 1. Mourning Dove | 55.58% frequency |
| 2. Red-winged Blackbird | 48.13% |
| 3. American Robin | 46.60% |
| 4. American Goldfinch | 42.15% |
| 5. Killdeer | 35.76% |
| 6. Eastern Kingbird | 35.20% |
| 7. Barn Swallow | 33.23% |
| 8. Western Meadowlark | 32.63% |
| 9. Common Grackle | 31.73% |
| 10. Song Sparrow | 30.59% |
| 11. Common Yellowthroat | 26.77% |
| 12. Chipping Sparrow | 26.63% |
| 13. Yellow Warbler | 25.75% |
| 14. Brown-headed Cowbird | 25.23% |
| 15. House Wren | 25.10% |
| 16. Western Kingbird | 24.60% |
| 17. Mallard | 22.92% |
| 18. House Sparrow | 21.42% |

| | |
|------------------------------|---------------|
| 19. Clay-colored Sparrow | 20.86% |
| 20. Ring-billed Gull | 19.77% |
| 21. Cliff Swallow | 19.38% |
| 22. Cedar Waxwing | 18.20% |
| 23. American Crow | 17.29% |
| 24. Northern Flicker | 17.18% |
| 25. Black-capped Chickadee | 16.69% |
| 26. Canada Goose | 16.68% |
| 27. American White Pelican | 15.97% |
| 28. Savannah Sparrow | 15.29% |
| 29. Yellow-headed Blackbird | 15.28% |
| 30. Red-tailed Hawk | 15.13% |
| 31. Marsh Wren | 14.81% |
| 32. Grasshopper Sparrow | 14.60% |
| 33. Tree Swallow | 14.57% |
| 34. Double-crested Cormorant | 14.50% |
| 35. Black Tern | 14.11% |
| 36. Blue-winged Teal | 14.10% |
| 37. Vesper Sparrow | 13.73% |
| 38. Horned Lark | 13.59% |
| 39. Least Flycatcher | 13.42% |
| 40. Gray Catbird | 13.09% |
| 41. Ring-necked Pheasant | 12.88% |

| | |
|-----------------------------|--------|
| 42. American Coot | 12.39% |
| 43. White-breasted Nuthatch | 12.22% |
| 44. Orchard Oriole | 11.99% |
| 45. Bobolink | 11.37% |
| 46. Gadwall | 11.26% |
| 47. House Finch | 10.89% |
| 48. Brewer's Blackbird | 10.68% |
| 49. Lesser Yellowlegs | 10.64% |
| 50. Franklin's Gull | 10.52% |
| 51. Ruddy Duck | 10.51% |
| 52. European Starling | 10.41% |
| 53. Lark Sparrow | 10.40% |
| 54. Spotted Towhee | 10.29% |
| 55. Upland Sandpiper | 10.28% |
| 56. Northern Harrier | 10.15% |
| 57. Bank Swallow | 9.73% |
| 58. Red-eyed Vireo | 9.66% |
| 59. Downy Woodpecker | 9.52% |
| 60. Spotted Sandpiper | 9.41% |
| 61. Turkey Vulture | 9.28% |
| 62. Sora | 9.25% |
| 63. Field Sparrow | 8.82% |
| 64. Eurasian Collared-Dove | 8.74% |

| | |
|-------------------------------|--------------|
| 65. Blue Jay | 8.42% |
| 66. Northern Shoveler | 8.40% |
| 67. Rock Pigeon | 8.21% |
| 68. Sedge Wren | 8.21% |
| 69. Lazuli Bunting | 8.05% |
| 70. Great Blue Heron | 7.78% |
| 71. Wilson's Phalarope | 7.72% |
| 72. Eared Grebe | 7.61% |
| 73. Swainson's Hawk | 7.55% |
| 74. Purple Martin | 7.54% |
| 75. Pied-billed Grebe | 7.47% |
| 76. Redhead | 7.27% |
| 77. Hairy Woodpecker | 7.26% |
| 78. Warbling Vireo | 7.21% |
| 79. American Avocet | 7.13% |
| 80. Greater Yellowlegs | 6.87% |
| 81. Brown Thrasher | 6.65% |
| 82. Western Grebe | 6.38% |
| 83. Wood Duck | 6.25% |
| 84. Baltimore Oriole | 6.00% |
| 85. Northern Pintail | 5.91% |
| 86. Forster's Tern | 5.87% |
| 87. American Redstart | 5.86% |

| | |
|---|--------------|
| 88. Wilson's Snipe | 5.86% |
| 89. Least Sandpiper | 5.83% |
| 90. American Kestrel | 5.83% |
| 91. Willet | 5.81% |
| 92. Dickcissel | 5.67% |
| 93. Yellow-breasted Chat | 5.48% |
| 94. Common Nighthawk | 5.42% |
| 95. Say's Phoebe | 5.25% |
| 96. Pectoral Sandpiper | 5.07% |
| 97. Chimney Swift | 4.97% |
| 98. Eastern Wood-Pewee | 4.90% |
| 99. Canvasback | 4.87% |
| 100. Lesser Scaup | 4.81% |
| 101. Chestnut-collared Longspur | 4.73% |
| 102. Black-billed Magpie | 4.54% |
| 103. Marbled Godwit | 4.43% |
| 104. Yellow-bellied Sapsucker | 4.41% |
| 105. White-faced Ibis | 4.23% |
| 106. Belted Kingfisher | 4.23% |
| 107. Bald Eagle | 4.19% |
| 108. Semipalmated Sandpiper | 4.14% |
| 109. Northern Rough-winged Swallow | 3.90% |
| 110. Green-winged Teal | 3.81% |

| | |
|--------------------------------|-------|
| 111. Willow Flycatcher | 3.77% |
| 112. Black-crowned Night Heron | 3.74% |
| 113. Great Egret | 3.71% |
| 114. Baird's Sandpiper | 3.58% |
| 115. Nelson's Sparrow | 3.45% |
| 116. Eastern Bluebird | 3.41% |
| 117. Sharp-tailed Grouse | 3.36% |
| 118. California Gull | 3.36% |
| 119. American Wigeon | 3.32% |
| 120. Stilt Sandpiper | 3.26% |
| 121. Great Crested Flycatcher | 3.20% |
| 122. Mountain Bluebird | 3.06% |
| 123. Cooper's Hawk | 2.99% |
| 124. Lark Bunting | 2.86% |
| 125. Red-headed Woodpecker | 2.73% |
| 126. Baird's Sparrow | 2.63% |
| 127. Solitary Sandpiper | 2.61% |
| 128. Hooded Merganser | 2.61% |
| 129. Rose-breasted Grosbeak | 2.55% |
| 130. Long-billed Dowitcher | 2.50% |
| 131. LeConte's Sparrow | 2.38% |
| 132. Red-necked Grebe | 2.27% |
| 133. Semipalmated Plover | 2.23% |

| | |
|---------------------------------------|--------------|
| 134. Black-headed Grosbeak | 2.23% |
| 135. American Bittern | 2.22% |
| 136. Wild Turkey | 2.20% |
| 137. Indigo Bunting | 2.17% |
| 138. Swamp Sparrow | 2.14% |
| 139. Rock Wren | 2.11% |
| 140. Ruby-throated Hummingbird | 2.04% |
| 141. Sprague's Pipit | 2.02% |
| 142. Eastern Phoebe | 2.01% |
| 143. Loggerhead Shrike | 1.92% |
| 144. Bufflehead | 1.86% |
| 145. Red-breasted Nuthatch | 1.84% |
| 146. Merlin | 1.81% |
| 147. Black-and-white Warbler | 1.80% |
| 148. Yellow-throated Vireo | 1.79% |
| 149. Ring-necked Duck | 1.77% |
| 150. Virginia Rail | 1.77% |
| 151. Common Tern | 1.76% |
| 152. Northern Cardinal | 1.71% |
| 153. Western Wood-Pewee | 1.68% |
| 154. Gray Partridge | 1.65% |
| 155. Red-necked Phalarope | 1.65% |
| 156. Short-billed Dowitcher | 1.60% |

| | |
|------------------------------------|----------------|
| 157. Great Horned Owl | 1.44% |
| 158. Western Cattle Egret | 1.25% |
| 159. Ovenbird | 1.21% |
| 160. Pine Siskin | 1.20% |
| 161. Golden Eagle | 1.18% |
| 162. Piping Plover | 1.13% |
| 163. Ferruginous Hawk | 1.10% |
| 164. Snowy Egret | 1.07% |
| 165. Veery | 1.07% |
| 166. Burrowing Owl | 1.06% |
| 167. Caspian Tern | 1.04% |
| 168. Common Raven | 1.02% |
| 169. Red-bellied Woodpecker | 0.9444% |
| 170. Sanderling | 0.9224% |
| 171. Black-billed Cuckoo | 0.8126% |
| 172. Green Heron | 0.7321% |
| 173. Violet-green Swallow | 0.7174% |
| 174. Least Tern | 0.7101% |
| 175. Pileated Woodpecker | 0.6955% |
| 176. Snow Goose | 0.6735% |
| 177. Horned Grebe | 0.6369% |
| 178. Common Loon | 0.5930% |
| 179. Prairie Falcon | 0.5783% |

| | |
|-------------------------------------|----------------|
| 180. Bullock's Oriole | 0.5783% |
| 181. Northern Waterthrush | 0.5051% |
| 182. Blue Grosbeak | 0.4978% |
| 183. Trumpeter Swan | 0.4832% |
| 184. Red Crossbill | 0.4612% |
| 185. Clark's Grebe | 0.4100% |
| 186. Least Bittern | 0.4100% |
| 187. Yellow Rail | 0.3807% |
| 188. Black-necked Stilt | 0.3733% |
| 189. Herring Gull | 0.3368% |
| 190. Buff-breasted Sandpiper | 0.3294% |
| 191. Short-eared Owl | 0.3294% |
| 192. Peregrine Falcon | 0.3221% |
| 193. Scarlet Tanager | 0.3221% |
| 194. Bonaparte's Gull | 0.3002% |
| 195. Hudsonian Godwit | 0.2855% |
| 196. Broad-winged Hawk | 0.2709% |
| 197. Osprey | 0.2562% |
| 198. Long-billed Curlew | 0.2489% |
| 199. Tennessee Warbler | 0.2489% |
| 200. Black-bellied Plover | 0.2343% |
| 201. Common Poorwill | 0.2123% |
| 202. Bell's Vireo | 0.1830% |

| | |
|---|----------------|
| 203. Eastern Towhee | 0.1684% |
| 204. Alder Flycatcher | 0.1611% |
| 205. Purple Finch | 0.1611% |
| 206. Common Goldeneye | 0.1464% |
| 207. Sandhill Crane | 0.1391% |
| 208. Ruddy Turnstone | 0.1391% |
| 209. Glossy Ibis | 0.1391% |
| 210. Brewer's Sparrow | 0.1391% |
| 211. Tundra Swan | 0.1318% |
| 212. American Black Duck | 0.1318% |
| 213. Sharp-shinned Hawk | 0.1318% |
| 214. Long-eared Owl | 0.1244% |
| 215. White-throated Sparrow | 0.1244% |
| 216. Neotropic Cormorant | 0.1025% |
| 217. Yellow-billed Cuckoo | 0.0952% |
| 218. Eastern Screech-Owl | 0.0952% |
| 219. Henslow's Sparrow | 0.0952% |
| 220. Yellow-rumped Warbler | 0.0952% |
| 221. Greater White-fronted Goose | 0.0878% |
| 222. Band-tailed Pigeon | 0.0878% |
| 223. Summer Tanager | 0.0805% |
| 224. Blue-gray Gnatcatcher | 0.0732% |
| 225. Eurasian Tree Sparrow | 0.0732% |

| | |
|-------------------------------------|----------------|
| 226. Ruffed Grouse | 0.0659% |
| 227. White-rumped Sandpiper | 0.0659% |
| 228. Thick-billed Longspur | 0.0659% |
| 229. Eastern Meadowlark | 0.0659% |
| 230. Orange-crowned Warbler | 0.0659% |
| 231. American Golden-Plover | 0.0586% |
| 232. Chestnut-sided Warbler | 0.0586% |
| 233. Common Merganser | 0.0512% |
| 234. Greater Prairie-Chicken | 0.0512% |
| 235. Red-throated Loon | 0.0439% |
| 236. Mourning Warbler | 0.0439% |
| 237. Cinnamon Teal | 0.0366% |
| 238. Eastern Whip-poor-will | 0.0366% |
| 239. Barred Owl | 0.0366% |
| 240. Northern Mockingbird | 0.0366% |
| 241. Nashville Warbler | 0.0366% |
| 242. Greater Sage-Grouse | 0.0293% |
| 243. Calliope Hummingbird | 0.0293% |
| 244. Rufous Hummingbird | 0.0293% |
| 245. American Woodcock | 0.0220% |
| 246. Dunlin | 0.0220% |
| 247. Brown Creeper | 0.0220% |
| 248. Red-breasted Merganser | 0.0146% |

| | | |
|------|-------------------------------------|----------------|
| 249. | White-winged Dove | 0.0146% |
| 250. | Red Knot | 0.0146% |
| 251. | Northern Saw-whet Owl | 0.0146% |
| 252. | Olive-sided Flycatcher | 0.0146% |
| 253. | Prairie Warbler | 0.0146% |
| 254. | Western Tanager | 0.0146% |
| 255. | Fulvous Whistling-Duck | 0.0073% |
| 256. | Ross's Goose | 0.0073% |
| 257. | Little Gull | 0.0073% |
| 258. | Great Black-backed Gull | 0.0073% |
| 259. | Lesser Black-backed Gull | 0.0073% |
| 260. | American Goshawk | 0.0073% |
| 261. | Yellow-bellied Flycatcher | 0.0073% |
| 262. | Scissor-tailed Flycatcher | 0.0073% |
| 263. | Blue-headed Vireo | 0.0073% |
| 264. | Wood Thrush | 0.0073% |
| 265. | American Pipit | 0.0073% |
| 266. | Dark-eyed Junco | 0.0073% |
| 267. | White-crowned Sparrow | 0.0073% |
| 268. | Blue-winged Warbler | 0.0073% |
| 269. | Black-bellied Whistling-Duck | 0.0000% |
| 270. | Greater Scaup | 0.0000% |
| 271. | Black Scoter | 0.0000% |

| | |
|--|--------------------------|
| 272. Whimbrel | 0.0000% |
| 273. Ruff | 0.0000% |
| 274. Little Stint | 0.0000% |
| 275. Western Sandpiper | 0.0000% |
| 276. South Polar Skua | 0.0000% |
| 277. Pacific Loon | 0.0000% |
| 278. Yellow-crowned Night Heron | 0.0000% |
| 279. Little Blue Heron | 0.0000% |
| 280. Rough-legged Hawk | 0.0000% |
| 281. Barn Owl | 0.0000% |
| 282. Snowy Owl | 0.0000% |
| 283. Crested Caracara | 0.0000% |
| 284. Swainson's Thrush | 0.0000% |
| 285. Hermit Thrush | 0.0000% |
| 286. Harris's Sparrow | 0.0000% |
| 287. Great-tailed Grackle | 0.0000% |
| 288. Cape May Warbler | 0.0000% |
| 289. Bay-breasted Warbler | 0.0000% |
| EXOTIC: ESCAPEE | |
| Mute Swan | 0.0659% frequency |
| Chukar | 0.0366% |
| Mandarin Duck | 0.0073% |
| African Collared-Dove | 0.0073% |

| | |
|--|--------------------------|
| Graylag Goose | 0.0000% |
| Northern Bobwhite | 0.0000% |
| HYBRIDS | |
| Lazuli x Indigo Bunting (hybrid) | 0.0512% frequency |
| Western x Clark's Grebe (hybrid) | 0.0439% |
| Spotted x Eastern Towhee (hybrid) | 0.0220% |
| Rose-breasted x Black-headed Grosbeak (hybrid) | 0.0146% |
| Snow x Ross's Goose (hybrid) | 0.0073% |
| Bullock's x Baltimore Oriole (hybrid) | 0.0073% |